



Town Center ORTHOPAEDICS

Baris Yildirim, MD
HAND & WRIST SURGERY
UPPER EXTREMITY SURGERY
1860 Town Center Drive, Ste 300
Reston, Virginia 20190
O: 703-435-6604 F: 703-662-4506

Post-operative Instructions: Distal Radius Fracture

- Start finger, elbow, and shoulder ROM immediately after surgery. Elevate as much as possible. No lifting/pushing/pulling with hand.
- Follow up with me 7-10 days after surgery. Go to hand therapist to change to removable wrist splint. OK to come out of splint for hygiene and daily ROM exercises. Start gentle active wrist ROM: flexion, extension, supination, pronation. Lift < 2lbs.
- Exceptions: if tenuous fixation, start wrist ROM at later date per surgeon.
- 2-6 weeks: Continue hand therapy with active wrist ROM exercises. Wear custom splint for activities and at night. Include intrinsic stretches. Continue finger, elbow, shoulder ROM until normal. Scar desensitization. Lift < 5 lbs.
- 6-12 weeks: Wean out of splint. Begin strengthening and passive ROM, lift < 10 lbs or 3 lbs if repetitive motion.
- Start work conditioning or return to sport program at 8 weeks.
- OK to return to full duty manual labor or contact sports at 3 months.